



UNIVERSITI PUTRA MALAYSIA

**STORAGE STABILITY OF CHILLED MINIMALLY PROCESSED
SHREDDED CABBAGE**

ROSHITA BINTI IBRAHIM

FSMB 2003 21

STORAGE STABILITY OF CHILLED MINIMALLY PROCESSED SHREDDED CABBAGE

By

ROSHITA BINTI IBRAHIM

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirement for the Degree of Master of Science**

September 2003



To my beloved family & friends... ..

Mak, Ayah, Kak Na, Kak Tie, Adik Nizam, Abang Amran, Abang Anwar, Abang Shah, Eja, Fanny, Ali, La, Kak Zie, Kak Norma, Ida, Anida, Kak Tie Kel, Kak Tie Mel, Nizam Lani, Fisal, Azam JB, Eed, Syafiq, Lynn Bio, Iza, Taufik and Along TJ.....

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirements for the degree of Master of Science

STORAGE STABILITY OF CHILLED MINIMALLY PROCESSED SHREDDED CABBAGE

By

ROSHITA BINTI IBRAHIM

July 2003

Chairman : Associate Professor Azizah Osman, Ph.D

Faculty : Food Science and Biotechnology

Minimally processed products are a rapidly growing segment for retail market and food service horticultural industry as they match the consumers' demand for product freshness and convenience. The shelf-life extension of these fresh-cut products is therefore relevant because of its economic impact. Packaging can be one of the important factors in determining the storage stability and quality of these fresh-cut products. The shelf life of minimally processed fruits and vegetables is often limited by enzymatic browning. The organoleptic properties of fruits and vegetables are strongly altered by the appearance of brown pigments. This study was conducted to determine the effects of using different types of polymeric films (Polypropylene (PP), Low Density Polyethylene (LDPE), High Density Polyethylene (HDPE) and PVC cling wrap (Control)) of varying permeabilities to gases and water vapour and also with and without the application of vacuum packaging on the physico-chemical, biochemical, microbiological characteristics and sensory aspects of the minimally processed (MP) shredded cabbage during storage at $5\pm 1^{\circ}\text{C}$; 90-95% RH. A study was

also conducted on the effects of dipping into 4 different anti-browning solutions namely 1% ascorbic acid, 0.1% sodium metabisulphite, 0.5% L-cysteine + 0.1% citric acid and 0.1% acetic acid on the storage stability of shredded cabbage. Sample dipped in distilled water was used as a control. Physico-chemical characteristics were determined by quantitative measurements of weight loss, colour, texture, ascorbic acid content, pH, titratable acidity, total soluble solid, chlorophyll content, polyphenol oxidase (PPO) activity and degree of browning. Carbon dioxide and ethylene production in the package atmosphere during storage were also determined with gas chromatography. The microbial characteristics determined were mesophilic and psychrotrophic bacterial counts and mold and yeast counts. Sensory evaluation involved subjective acceptability and descriptive analyses. Data collected were analyzed using ANOVA and Duncan Multiple Range Test (DMRT) at 5% significant level. Generally, in almost all the analyses done, the quality of the MP shredded cabbage deteriorated with increase in storage time. Among all the packaging films used, it was found that PP which is the least permeable film for gases and water vapour, could extend the shelf life of the MP shredded cabbage almost up to 3 weeks with minimum colour change, reduction in ascorbic acid content and deterioration in sensory properties, and marginally low changes in other parameters tested. Whilst PVC cling wrap (control) was found to be the least effective packaging film. Different packaging systems did not affect the microflora of the shredded cabbage which was predominantly bacteria, small numbers of yeasts and only an occasional mold for both mesophilic and psychrotrophic microorganisms. Samples packed in vacuum packaging showed no significance difference with those in non-vacuum packaging in almost all the parameters tested for all the different packaging films, even though the air from the package headspace which can cause oxidation spoilage to the produce

had been removed. Anti-browning treatment of 0.1% sodium metabisulphite gave the best sensory properties and visual colour retention followed by 0.1% acetic acid solution. Mean while 0.5% L-cysteine + 0.1% citric acid and 1% ascorbic acid solutions were found to be not very good anti-browning agents for the MP shredded cabbage as they gave worse results compared to control.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

KESTABILAN PENYIMPANAN DINGIN HIRISAN KOBIS TERPROSES MINIMA

Oleh

ROSHITA BINTI IBRAHIM

Julai 2003

Pengerusi : Professor Madya Azizah Osman, Ph.D.

Fakulti : Sains Makanan dan Bioteknologi

Produk-produk pemprosesan minima berkembang pesat di dalam pasaran jualan runcit dan industri perkhidmatan makanan berasaskan produk hortikultur kerana ia memenuhi permintaan pengguna terhadap produk yang segar dan mudah penyediaannya. Pemanjangan hayat penyimpanan produk-produk potong-segar ini adalah penting kerana penjimatannya dari segi ekonomi. Pembungkusan boleh menjadi salah satu daripada faktor penting dalam mengekalkan hayat penyimpanan produk-produk pemprosesan minima ini. Hayat penyimpanan pemprosesan minima buah-buahan dan sayur-sayuran selalunya terhad disebabkan oleh masalah keperangan akibat aktiviti enzim. Ciri-ciri organoleptiknya juga berubah dengan wujudnya kesan keperangan pada produk itu. Kajian ini telah dijalankan untuk menentukan kesan-kesan yang didapati dengan penggunaan filem-filem plastik yang berbeza untuk pembungkusan iaitu, Polipropilena (PP), Polietilina berketumpatan rendah (LDPE), Polietilina berketumpatan tinggi (HDPE) dan selaput nipis Poli vinilklorida (PVC)(kawalan) yang mempunyai ketelapan yang berbeza terhadap gas dan wap air

serta dengan pengaplikasian pembungkusan vakum dan tanpa vakum terhadap ciri-ciri fisiko-kimia, biokimia, mikrobiologi dan aspek-aspek deria ke atas pemprosesan minimal hirisan kobis semasa penyimpanan pada suhu $5\pm 1^{\circ}\text{C}$ dan kelembapan relatif 90-95%. Kajian terhadap kesan-kesan penggunaan larutan penyahperang yang berlainan terhadap hayat penyimpanan hirisan kobis juga dijalankan. Larutan-larutan penyahperang yang digunakan ialah 1% asid askorbik, 0.1% natrium metabisulfite, campuran 0.5% L-cysteine dan 0.1% asid sitrik dan 0.1% asid asetik. Air suling digunakan sebagai larutan kawalan. Ciri-ciri fisiko-kimia yang ditentukan secara kuantitatif merangkumi ujian-ujian seperti % kehilangan berat sampel, perubahan warna, tekstur, kandungan asid askorbik, pH, keasidan titratan, jumlah pepejal larut, kandungan klorofil, aktiviti enzim polyphenol oxidase (PPO) dan tahap keperangan sampel. Penghasilan gas karbon dioksida dan etilina di dalam bungkusan sampel semasa penyimpanan juga ditentukan dengan kromatografi gas. Analisa mikrob dilakukan untuk melihat kehadiran bakteria mesofilik dan psikrotropik serta kulat dan yis. Penilaian deria pula dilakukan berdasarkan penerimaan subjektif dan analisa perubahan rupa bentuk sampel. Data dianalisa dengan ANOVA dan Duncan Multiple Range Test (DMRT) pada tahap signifikan 5%. Secara umumnya, dalam hampir semua ujian yang dilakukan kualiti hirisan kobis semakin menurun dengan bertambahnya masa penyimpanan. Antara semua filem-filem plastik pembungkus yang digunakan, PP merupakan filem yang paling tidak telap kepada gas dan wap air, boleh memanjangkan hayat penyimpanan hirisan kobis kepada hampir 3 minggu dengan perubahan warna, pengurangan kandungan asid askorbik dan pengurangan ciri-ciri deria yang minimal serta sedikit perubahan dalam ujian-ujian lainnya. Sementara selaput nipis PVC (kawalan) merupakan filem yang paling tidak efektif. Kaedah pembungkusan yang berbeza tidak menunjukkan perbezaan yang bererti

kepada pertumbuhan mikrob di dalam hirisan kobis yang kebanyakannya terdiri daripada bakteria, sedikit yis serta kulat untuk kedua-dua jenis mesofilik dan psikrotropik. Walaupun udara di dalam bungkusan telah dinyahkan, pembungkusan vakum tidak menunjukkan perbezaan yang bererti dengan pembungkusan tanpa vakum dalam hampir kesemua parameter-parameter yang diuji untuk berlainan jenis filem-filem plastik yang digunakan. Larutan penyahperang 0.1% natrium metabisulfite memberikan keputusan yang terbaik dari segi ciri-ciri deria dan pengekalan warna sampel, diikuti oleh larutan 0.1% asid asetik. Manakala larutan campuran 0.5% L-cysteine dan 0.1% asid sitrik dan larutan 1% asid askorbik tidak boleh menjadi agen-agen penyahperang yang baik bagi hirisan kobis memandangkan larutan-larutan ini memberikan keputusan yang lebih buruk daripada larutan kawalan.

ACKNOWLEDGEMENTS

First and foremost, Alhamdulillah, my utmost thanks and gratitude to the Almighty Allah S.W.T, the Sustainer, the Most Gracious and Most Merciful, who has given me the strength and capability to complete this Master Degree, and my salawat and salam to His righteous messenger, prophet Muhammad S.A.W.

Sincerest gratitude and appreciation to the chairman of my Supervisory Committee, Assoc. Prof. Dr. Azizah Osman and other members of my supervisory committee, Assoc. Prof. Dr. Russly Abdul Rahman and Assoc. Prof. Dr. Nazamid Saari for their unwavering support, invaluable guidance, suggestions, discussion and patience throughout the project and also for their constructive comments and criticisms towards the research and preparation of this thesis.

My sincere gratitude is also extended to MOSTE for the financial support, through IRPA Grant No: 01-02-04-0519 which is awarded to Assoc. Prof. Dr. Azizah Osman to conduct this study. My heartfelt thanks also go to all the dedicated staffs of Food Science and Food Technology Departments and also to the graduate students in Postharvest Lab, Hazniza, Fanny, Ali, Dwi, Taufiq and Iza for their generous cooperation, assistance, guidance and support. Acknowledgements is also due to all my housemates and friends in Fish Lab, Nutraceutical Lab, Microbiology Lab and also all my very best friends in KUSTEM who have given me the moral encouragement and support to complete my study. It is quite impossible to list all their names here. To my

head of department, Pn. Wan Sarah Wan Abdullah and all the lecturers at JSM, KUSTEM, thank you very much for their consideration, helps and understanding. I would also like to extend my thanks to Marditech Plantation, Cameron Highland for their cooperation.

Finally, I also wish to express my deepest appreciation and love to my parents, sisters and brothers who have been very patience and given me encouragement and support during the many years of my seemingly never ending pursue for knowledge. I wish for every bead of sweat produced will be in Allah's barakah.

I certify that an Examination Committee met on 24th July 2003 to conduct the final examination of Roshita binti Ibrahim on her Master of Science thesis entitled “Storage Stability of Chilled Minimally Processed Shredded Cabbage” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

SALMAH YUSOF, Ph.D.

Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

AZIZAH OSMAN, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

NAZAMID SAARI, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

RUSSLY ABDUL RAHMAN, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor / Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 30 SEP 2003

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows:

AZIZAH OSMAN, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

NAZAMID SAARI, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

RUSSLY ABDUL RAHMAN, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph.D.
Professor / Dean
School of Graduate Studies
Universiti Putra Malaysia

Date : **14** NOV 2003

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.


ROSHITA IBRAHIM

Date : 28/9/03

TABLE OF CONTENTS

DEDICATION.....	Page ii
ABSTRACT.....	iii
ABSTRAK.....	vi
ACKNOWLEDGEMENTS.....	ix
APPROVAL.....	xi
DECLARATION.....	xiii
LIST OF TABLES.....	xvii
LIST OF FIGURES.....	xx
LIST OF PLATES.....	xxiv
LIST OF ABBREVIATIONS.....	xxvi

CHAPTER

1	GENERAL INTRODUCTION.....	1
2	LITERATURE REVIEW.....	7
2.1	Introduction to Cabbage.....	7
2.2	Cabbage Production in Cameron Highland, Malaysia.....	9
2.3	Postharvest Handling of Cabbage.....	10
2.4	Minimally Processed Horticultural Products.....	12
2.5	Potential Degradation in Minimally Processed Products.....	13
2.5.1	Respiration Rate.....	13
2.5.2	Microbial Spoilage.....	14
2.5.3	Textural Changes.....	15
2.5.4	Nutrients Loss.....	16
2.5.5	Colour Changes (Browning).....	17
2.5.5.1	Degradation of Chlorophyll.....	17
2.5.5.2	Oxidative Reaction of Phenolic Compound.....	19
2.5.5.3	Polyphenol oxidase.....	20
2.6	Factors Affecting Storage Stability of Minimally Processed Fruits and Vegetables.....	21
2.6.1	Temperature.....	21
2.6.2	Relative Humidity.....	22
2.6.3	Microorganisms.....	23
2.6.4	Packaging.....	24
2.6.4.1	Forms of Packaging.....	25
2.6.4.2	Packaging Materials.....	27
2.6.4.3	Effects of Packaging on the Shelf life of Fresh Produce.....	31
2.6.5	Anti-browning Treatments.....	32
2.6.5.1	Physical Preservatives.....	33
2.6.5.2	Chemical Preservatives.....	33

2.6.6	Others.....	37
3	EFFECTS OF DIFFERENT PACKAGING FILMS ON THE PHYSICO-CHEMICAL CHARACTERISTICS AND SENSORY ASPECTS OF SHREDDED CABBAGE STORED AT CHILLED TEMPERATURE WITH AND WITHOUT THE APPLICATION OF VACUUM.....	39
3.1	Introduction.....	39
3.2	Materials and Methods.....	40
3.2.1	Source of Cabbage Sample.....	40
3.2.2	Packaging Materials.....	40
3.2.3	Sample Preparation.....	42
3.2.4	Experimental Design.....	42
3.2.5	Physical Analyses.....	43
3.2.5.1	Percentage of Physiological Weight Loss.....	43
3.2.5.2	Determination of Colour.....	43
3.2.5.3	Texture Analysis.....	45
3.2.6	Chemical Analyses.....	45
3.2.6.1	Determination of Ascorbic Acid Content.....	45
3.2.6.2	Determination of pH.....	45
3.2.6.3	Determination of Titratable Acidity.....	46
3.2.6.4	Determination of Total Soluble Solid (TSS).....	46
3.2.7	Sensory Evaluation.....	46
3.2.8	Statistical Analysis.....	46
3.3	Results and Discussion.....	47
3.3.1	Physical Changes.....	47
3.3.1.1	Percentage Physiological Weight Loss.....	47
3.3.1.2	Colour.....	51
3.3.1.3	Texture.....	57
3.3.2	Chemical Changes.....	63
3.3.2.1	Ascorbic Acid Content.....	63
3.3.2.2	pH.....	69
3.3.2.3	Titratable Acidity.....	71
3.3.2.4	Total Soluble Solid.....	73
3.3.3	Sensory Evaluation.....	75
3.4	Conclusion.....	84
4	EFFECTS OF DIFFERENT PACKAGING FILMS ON THE CO₂ AND C₂H₄ PRODUCTION, PPO ACTIVITY, DEGREE OF BROWNING, CHLOROPHYLL CONTENT AND MICROBIOLOGICAL STUDY ON THE SHREDDED CABBAGE STORED AT CHILLED TEMPERATURE WITH AND WITHOUT THE APPLICATION OF VACUUM.....	85
4.1	Introduction.....	85
4.2	Materials and Methods.....	87
4.2.1	Determination of Ethylene and CO ₂ Production in the Package Atmosphere.....	88
4.2.2	Determination of PPO Activity.....	89
4.2.2.1	Preparation of Shredded Cabbage for PPO Extract...	89
4.2.2.2	Assay for PPO Activity.....	89

4.2.3	Determination of Degree of Browning.....	89
4.2.4	Determination of Chlorophyll Content.....	90
4.2.5	Microbiological Study.....	90
4.3	Results and Discussion.....	91
4.3.1	CO ₂ and C ₂ H ₄ Production in the Package Atmosphere.....	91
4.3.2	PPO Activity.....	95
4.3.3	Degree of Browning.....	100
4.3.4	Chlorophyll Content.....	102
4.3.5	Microbiological Study.....	104
4.3.5.1	Mesophilic Microorganisms.....	104
4.3.5.2	Psychrotrophic Microorganisms.....	111
4.4	Conclusion.....	115
5	EFFECTS OF SELECTED ANTI-BROWNING TREATMENT ON THE STORAGE QUALITY OF SHREDDED CABBAGE.....	116
5.1	Introduction.....	116
5.2	Materials and Methods.....	118
5.2.1	Anti-browning Agents.....	120
5.2.2	Determination of Colour Changes.....	120
5.2.3	Sensory Evaluation.....	120
5.3	Results and Discussion.....	121
5.3.1	Colour (Hunter Lab).....	121
5.3.2	Sensory Evaluation.....	125
5.4	Conclusion.....	127
6	CONCLUSIONS AND RECOMMENDATIONS	131
6.1	Conclusions.....	131
6.2	Recommendations.....	132
	BIBLIOGRAPHY.....	133
	APPENDICES.....	145
A1	Determination of Titratable Acidity and pH.....	146
A2	Determination of Vitamin C.....	147
A3	Texture Analyses.....	148
A4	Microbiological Study.....	149
A5	Sensory Evaluation.....	151
B	Additional Figures.....	155
C	Additional Tables.....	171
D	Plates.....	192
	BIODATA OF THE AUTHOR/VITA.....	207

LIST OF TABLES

Table	Page
1 Nutritional Content of 1cup (69 g) of Shredded Cabbage.....	8
2 Overall mean values for Physiological Weight Loss and Colour Changes (Hunter L*, a* and b* values) of Minimally Processed Shredded Cabbage Packed in Different Packaging Films with Vacuum and Non-vacuum Conditions During Storage.....	48
3 Overall mean values for Texture Analysis (Firmness and Crispness of Leaves and Midrib) of Minimally Processed Shredded Cabbage Packed in Different Packaging Films with Vacuum and Non-vacuum Conditions During Storage.....	58
4 Overall mean values for Ascorbic Acid Content, pH, Titratable Acidity and TSS of Minimally Processed Shredded Cabbage Packed in Different Packaging Films with Vacuum and Non-vacuum Conditions During Storage.....	66
5 Overall mean values for Sensory Evaluation (Hedonic) of Minimally Processed Shredded Cabbage Packed in Different Packaging Films During Storage.....	77
6 Overall mean values for Sensory Evaluation (QDA) of Minimally Processed Shredded Cabbage Packed in Different Packaging Films During Storage.....	80
7 Overall mean values for CO ₂ and C ₂ H ₄ Production in the Package Atmosphere of Minimally Processed Shredded Cabbage Packed in Different Packaging Films Vacuum and Non-vacuum Conditions During Storage.....	92
8 Overall mean values for PPO Activity, Degree of Browning and Chlorophyll Content of Minimally Processed Shredded Cabbage Packed in Different Packaging Films with Vacuum and Non-vacuum Conditions During Storage.....	97
9 Overall mean values for Microbiology Study (Mesophilic and Psychrotrophic Bacterial, Yeast and Mold Count) of Minimally Processed Shredded Cabbage Packed in Different Packaging Films with Vacuum and Non-vacuum Conditions During Storage.....	107

10	Overall mean values for Colour (Hunter L*, a* & b* value) of Minimally Processed Shredded Cabbage Dipped in Anti-browning Solutions During Storage.....	123
11	Overall mean values for Sensory Evaluation (Hedonic) of Minimally Processed Shredded Cabbage Dipped in Different Anti-browning Solutions During Storage.....	126
12	Overall mean values for Sensory Evaluation (QDA) of Minimally Processed Shredded Cabbage Dipped in Different Anti-browning Solutions During Storage.....	128
13	Effects of Vacuum and Different Packaging Films on the Physiological Weight Loss and Colour Changes (Hunter L*, a* and b* values) of Minimally Processed Shredded Cabbage During Storage.....	171
14	Effects of Vacuum and Different Packaging Films on the Texture (Firmness and Crispness of Leaves and Midrib) of Minimally Processed Shredded Cabbage During Storage.....	173
15	Effects of Vacuum and Different Packaging Films on the Ascorbic Acid Content, pH, Titratable Acidity and TSS of Minimally Processed Shredded Cabbage During Storage.....	175
16	Effects of Vacuum and Different Packaging Films on the Sensory Evaluation (Hedonic) of Minimally Processed Shredded Cabbage During Storage.....	177
17	Effects of Vacuum and Different Packaging Films on the Sensory Evaluation (QDA) of Minimally Processed Shredded Cabbage During Storage.....	179
18	Effects of Vacuum and Different Packaging Films on the CO ₂ and C ₂ H ₄ Production in the Package Atmosphere of Minimally Processed Shredded Cabbage During Storage.....	181
19	Effects of Vacuum and Different Packaging Films on the PPO Activity, Degree of Browning and Chlorophyll Content of Minimally Processed Shredded Cabbage During Storage.....	182
20	Effects of Vacuum and Different Packaging Films on the Microbial Study (Mesophilic and Psychrotrophic Bacterial, Yeast and Mold Count) of Minimally Processed Shredded Cabbage During Storage.....	184
21	Effects of Dipping into Anti-browning Solutions on the Colour (Hunter L*, a* & b* value) Changes of Minimally Processed Shredded Cabbage During Storage.....	186
22	Effects of Dipping into Anti-browning Solutions on the Sensory Evaluation (Hedonic) of the Minimally Processed Shredded Cabbage	

	During Storage.....	188
23	Effects of Dipping into Anti-browning Solutions on the Sensory Evaluation (QDA) of the Minimally Processed Shredded Cabbage During Storage.....	190

LIST OF FIGURES

Figure		Page
1	Pathways of Chlorophyll Degradation in Green Tissue.....	18
2	Oxidation Reaction in Enzymatic Browning.....	21
3	Flow Diagram on the Sample Preparation and Processes.....	44
4	Effect of Different Packaging Films on the % Physiological Weight Loss of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	50
5	Effect of Different Packaging Films on the Colour Changes (Hunter L* Values) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	52
6	Effect of Different Packaging Films on the Colour Changes (Hunter a* Values) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	54
7	Effect of Different Packaging Films on the Colour Changes (Hunter b* Values) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	55
8	Effect of Different Packaging Films on the Firmness (Compression Force) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	60
9	Effect of Different Packaging Films on the Leaves Crispness (Shearing Force) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	62
10	Effect of Different Packaging Films on the Midribs Crispness (Shearing Force) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	64

11	Effect of Different Packaging Films on the Ascorbic acid Content of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	68
12	Effect of Different Packaging Films on the pH of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	70
13	Effect of Different Packaging Films on the Titratable Acidity (% Malic acid) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	72
14	Effect of Different Packaging Films on the Total Soluble Solid (°Brix) of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	74
15	Sensory Evaluation (QDA) of Minimally Processed Shredded Cabbage Packed in Different Packaging Films (a) Non-vacuumed (b) Vacuumed During Low Temperature Storage.....	83
16	Effect of Different Packaging Films on the (a) Carbon dioxide and (b) Ethylene Production in the Package Atmosphere of Minimally Processed Shredded Cabbage During Low Temperature Storage.....	94
17	Effect of Different Packaging Films on the Polyphenol Oxidase (PPO) Activity of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	99
18	Effect of Different Packaging Films on the Degree of Browning of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	101
19	Effect of Different Packaging Films on the Chlorophyll Content of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	103
20	Effect of Different Packaging Films on the Mesophilic Bacteria Count of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	109
21	Effects of Different Packaging Films on the Mesophilic Yeast and Mold Count of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage.....	110
22	Effects of Different Packaging Films on the Psychrotrophic Bacteria Count of Minimally Processed Shredded Cabbage. (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature	

	Storage	112
23	Effects of Different Packaging Films on the Psychrotrophic Yeast and Mold Count of Minimally Processed Shredded Cabbage (a) Non-vacuum Packaging and (b) Vacuum Packaging During Low Temperature Storage	114
24	Flow Diagram on the Sample Preparation and Processes (Anti-browning Treatment)	119
25	Colour Changes (a) Hunter L*, (b) Hunter a* and (c) Hunter b* Values of Minimally Processed Shredded Cabbage Dipped in Different Anti-browning Solutions During Low Temperature Storage	124
26	Sensory Evaluation (QDA) of Minimally Processed Shredded Cabbage Dipped in Different Anti-browning Solutions During Low Temperature Storage	129
27a	Effect of Different Packaging Films on the Sensory Acceptability (Colour) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	155
27b	Effect of Different Packaging Films on the Sensory Acceptability (Texture) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	156
27c	Effect of Different Packaging Films on the Sensory Acceptability (Odour) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	157
27d	Effect of Different Packaging Films on the Sensory Acceptability (Freshness) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	158
27e	Effect of Different Packaging Films on the Sensory Acceptability (Overall Acceptability) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	159
28a	Effect of Different Packaging Films on the Sensory (QDA-Yellowish White) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	160
28b	Effect of Different Packaging Films on the Sensory (QDA-Brightness) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	161
28c	Effect of Different Packaging Films on the Sensory (QDA-Texture) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed	162

28d	Effect of Different Packaging Films on the Sensory (QDA-Odour) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed.....	163
28e	Effect of Different Packaging Films on the Sensory (QDA-Freshness) of Minimally Processed Shredded Cabbage During Storage (i) Non-vacuumed (ii) Vacuumed.....	164
29	Effects of Dipping into Anti-browning Solutions on the Sensory Acceptability (Hedonic) of the Minimally Processed Shredded Cabbage During Storage (a) Colour Acceptability, (b) Texture Acceptability, (c) Odour Acceptability, (d) Freshness Acceptability and (e) Overall Acceptability.....	165
30	Effects of Dipping into Anti-browning Solutions on the Sensory Evaluation (QDA) of the Minimally Processed Shredded Cabbage During Storage (a) Yellowish White, (b) Brightness, (c) Texture, (d) Odour and (e) Freshness.....	168

LIST OF PLATES

Plate		Page
1	Marditech Cabbage Plantation in Cameron Highland.....	192
2	Head of Cabbage Reaching Maturity.....	192
3	National Food Processor (MK 5070M) – Shredder.....	193
4	Removing Outer Leaves and Core.....	193
5	Cabbage: Before and After Shredding.....	194
6	Washing/Dipping Shredded Cabbage in Water for 30 sec.....	194
7	Toast-drying Shredded Cabbage in Basket after Washing.....	195
8	400g of Shredded Cabbage Packed in Different Packaging Films (Day 0)	195
9	MP Shredded Cabbage Packed in Different Packaging Films after 1 week of Storage.....	196
10	MP Shredded Cabbage Packed in Different Packaging Films after 2 week of Storage.....	196
11	MP Shredded Cabbage Packed in Different Packaging Films after 3 week of Storage.....	197
12	MP Shredded Cabbage Packed in Different Packaging Films after 4 week of Storage.....	197
13	MP Shredded Cabbage were Vacuum-packed in Different Packaging Films (Day 0).....	198
14	MP Shredded Cabbage were Vacuum-packed in Different Packaging Films after 1 week of Storage.....	198
15	MP Shredded Cabbage were Vacuum-packed in Different Packaging Films after 2 week of Storage.....	199
16	MP Shredded Cabbage were Vacuum-packed in Different Packaging Films after 3 week of storage.....	199